



**IMMUNIZE
WISCONSIN**

CLINICIAN GUIDANCE:

COVID-19 DOSING CHARTS

Last Updated December 4, 2024

The information listed above was developed in coordination with Illinois Chapter, American Academy of Pediatrics via the Centers for Disease Control and Prevention vaccination administration guidelines.

COVID-19 VACCINATION SCHEDULE AND DOSING

AGES 6 MONTHS TO 4 YEARS




UNVACCINATED

dose/injection volume

Moderna 2024-2025:
(Do NOT dilute before use)
Dark Blue Cap (green label)

Pfizer 2024-2025:
(dilute before use)
Yellow Cap




PREVIOUSLY VACCINATED

dose/injection volume

☐ **Previously Received COVID-19 Vaccines**

Moderna 2024-2025:
(Do NOT dilute before use)
Dark Blue Cap (green label)

Pfizer 2024-2025:
(dilute before use)
Yellow Cap



PLEASE NOTE

An 8-week interval between the first and second COVID-19 vaccine (Moderna and Pfizer-BioNTech) doses might be optimal for some people as it might reduce the rare risk of myocarditis and pericarditis associated with these vaccines.

The information listed above was developed in coordination with Illinois Chapter, American Academy of Pediatrics via the Centers for Disease Control and Prevention vaccination administration guidelines.

immunizewi.org

COVID-19 VACCINATION SCHEDULE AND DOSING

AGES 6 MONTHS TO 4 YEARS IMMUNOCOMPROMISED




UNVACCINATED

dose/injection volume

Moderna 2024-2025:
(Do NOT dilute before use)
Dark Blue Cap (green label)

Pfizer 2024-2025:
Yellow Cap (dilute before use)




PREVIOUSLY VACCINATED

dose/injection volume

☐ **Previously Received COVID-19 Vaccines**

Moderna 2024-2025:
(Do NOT dilute before use)
Dark Blue Cap (green label)

Pfizer 2024-2025:
Yellow Cap (dilute before use)



ADDITIONAL DOSES

Further additional 2024-2025 mRNA dose(s) may be administered from the same manufacturer, informed by the clinical judgment of a healthcare provider and personal preference and circumstances. Additional doses should be administered at least 2 months after the last 2024-2025 mRNA vaccine dose.

The information listed above was developed in coordination with Illinois Chapter, American Academy of Pediatrics via the Centers for Disease Control and Prevention vaccination administration guidelines.

immunizewi.org

COVID-19 VACCINATION SCHEDULE AND DOSING

AGES 5 TO 11 YEARS



UNVACCINATED

dose/injection volume

Moderna 2024-2025:


(Do NOT dilute before use)

Dark Blue Cap (green label)

Pfizer 2024-2025:

(Do NOT dilute before use)

Blue Cap



PREVIOUSLY VACCINATED

dose/injection volume



**Previously Received
COVID-19 Vaccines**

Moderna 2024-2025:


(Do NOT dilute before use)

Dark Blue Cap (green label)

Pfizer 2024-2025:

(Do NOT dilute before use)

Blue Cap



The information listed above was developed in coordination with Illinois Chapter, American Academy of Pediatrics via the Centers for Disease Control and Prevention vaccination administration guidelines.

immunizewi.org

COVID-19 VACCINATION SCHEDULE AND DOSING

AGES 5 TO 11 YEARS IMMUNOCOMPROMISED




UNVACCINATED

dose/injection volume

Moderna 2024-2025:
(Do NOT dilute before use)
Dark Blue Cap (green label)

Pfizer 2024-2025:
Blue Cap (Do NOT dilute before use)




PREVIOUSLY VACCINATED

dose/injection volume

☐ **Previously Received COVID-19 Vaccines**

Moderna 2024-2025:
(Do NOT dilute before use)
Dark Blue Cap (green label)

Pfizer 2024-2025:
Blue Cap (Do NOT dilute before use)



Layered vaccine manufacturer instances indicate **interchangeable doses**.

ADDITIONAL DOSES

Further additional 2024-2025 mRNA dose(s) may be administered, informed by the clinical judgment of a healthcare provider and personal preference and circumstances. Additional doses should be administered at least 2 months after the last 2024-2025 mRNA vaccine dose.

The information listed above was developed in coordination with Illinois Chapter, American Academy of Pediatrics via the Centers for Disease Control and Prevention vaccination administration guidelines.

immunizewi.org

COVID-19 VACCINATION SCHEDULE AND DOSING

AGES 12 YEARS AND OLDER




UNVACCINATED

dose/injection volume

Moderna 2024-2025:
(Do NOT dilute before use)
Dark Blue Cap (dark blue label)

Pfizer 2024-2025:
(Do NOT dilute before use)
Gray Cap




PREVIOUSLY VACCINATED

dose/injection volume

☐ **Previously Received COVID-19 Vaccines**

Moderna 2024-2025:
(Do NOT dilute before use)
Dark Blue Cap (dark blue label)

Pfizer 2024-2025:
(Do NOT dilute before use)
Gray Cap



The information listed above was developed in coordination with Illinois Chapter, American Academy of Pediatrics via the Centers for Disease Control and Prevention vaccination administration guidelines.

immunizewi.org

COVID-19 VACCINATION SCHEDULE AND DOSING

AGES 12 YEARS AND OLDER IMMUNOCOMPROMISED




UNVACCINATED

dose/injection volume

Moderna 2024-2025:
(Do NOT dilute before use)
Dark Blue Cap (dark blue label)

Pfizer 2024-2025:
(Do NOT dilute before use)
Gray Cap




PREVIOUSLY VACCINATED

dose/injection volume

☐ **Previously Received COVID-19 Vaccines**

Moderna 2024-2025:
(Do NOT dilute before use)
Dark Blue Cap (dark blue label)

Pfizer 2024-2025:
(Do NOT dilute before use)
Gray Cap



Layered vaccine manufacturer instances indicate **interchangeable doses.**

ADDITIONAL DOSES

Further additional 2024-2025 dose(s) may be administered, informed by the clinical judgment of a healthcare provider and personal preference and circumstances. Additional doses should be administered at least 2 months after the last 2024-2025 vaccine dose.

The information listed above was developed in coordination with Illinois Chapter, American Academy of Pediatrics via the Centers for Disease Control and Prevention vaccination administration guidelines.

immunizewi.org

COVID-19 VACCINATION SCHEDULE AND DOSING


AGES 12 YEARS AND OLDER



UNVACCINATED

dose/injection volume

Novavax 2024-2025:
(Do NOT dilute before use)
Royal Blue Cap





PREVIOUSLY VACCINATED

dose/injection volume



Previously Received
COVID-19 Vaccines

Novavax 2024-2025:
(Do NOT dilute before use)
Royal Blue Cap



PLEASE NOTE

An 8-week interval between the first and second COVID-19 vaccine doses might be optimal for some people as it might reduce the rare risk of myocarditis and pericarditis associated with these vaccines.

The information listed above was developed in coordination with Illinois Chapter, American Academy of Pediatrics via the Centers for Disease Control and Prevention vaccination administration guidelines.

immunizewi.org

COVID-19 VACCINATION SCHEDULE AND DOSING

AGES 12 YEARS AND OLDER IMMUNOCOMPROMISED



UNVACCINATED

dose/injection volume

Novavax 2024-2025:
(Do NOT dilute before use)
Royal Blue Cap



PREVIOUSLY VACCINATED

dose/injection volume

☐ Previously Received
COVID-19 Vaccines

Novavax 2024-2025:
(Do NOT dilute before use)
Royal Blue Cap



Layered vaccine
manufacturer
instances indicate
interchangeable doses.

ADDITIONAL DOSES

Further additional 2024-2025 dose(s) may be administered, informed by the clinical judgment of a healthcare provider and personal preference and circumstances. Additional doses should be administered at least 2 months after the last 2024-2025 vaccine dose.

The information listed above was developed in coordination with Illinois Chapter, American Academy of Pediatrics via the Centers for Disease Control and Prevention vaccination administration guidelines.

immunizewi.org